

# Myna Bird Trapping Help Sheet

January 2015

This help sheet is intended as a guide to trapping Common Mynas aka Indian Mynas. It has been prepared with help from CIMAC, the Canberra action Committee. **Traps are to be placed only on private property, to avoid possible public sensitivity and public liability issues and to avoid vandalism.**

- Place the trap in a relatively open area or where birds already feed. A good spot seems to be under power lines or in an area where there are not people or animal traffic (ie behind the carport or in the vegie patch).
- Bait the trap with white bread or a food the birds are accustomed to feeding on. Suburban feeding birds are used to pet food as many pet owners leave food out for their pets during the day. Using small dry dog and cat biscuits has been very successful as has VIP pet roll. We generally do not use cereal and grain-based foods (birdseed etc) as this attracts non-target birds such as crested pigeons and parrots. However, slices or pieces of white bread work really well in North Queensland.
- To attract the Mynas to the area, put a little loose feed outside the funnel entrances, in the tunnel, and **under** the feeding chamber where the Mynas can see the food directly in front of them, when they are in the tunnel. Put food in the collection box for captured birds, and provide fresh water.
- Loose feeding will attract the birds into your area. You should not over feed outside the trap, as this will not encourage the birds to enter the trap. Use loose feed only to encourage the birds to feed, not to feed them.
- Monitor daily to ensure bait and water are available while trapping.
- Keep pets (particularly cats) away when trapping as the birds see cats and dogs as predators and are unlikely to approach the trap.
- Clean (hose down) and relocate the trap after the area becomes soiled from captured birds, as they don't appear to like being around their own excrement. Place newspaper or cardboard under the trap if using the trap on paved surfaces to avoid spoiling your pavements.
- Don't be disappointed if you don't catch birds every day. They are spasmodic in their movements, so keep at it. They may have moved to another food source in the area but will return if you keep feeding.
- If you keep a call bird in the containment chamber of your trap, this is likely to encourage others to the trap. This is not essential, and if you do this you must provide captive birds with food, water and shelter.
- Use gloves when handling live or dead birds as wild birds may carry disease.
- Abide by the Cairns RIM Protocol on Animal Welfare: this has been cleared by the RSPCA.
- If you not in a position to euthanize the captured birds yourself, please approach the Earlville Veterinarian Surgery at 474 Mulgrave Rd for euthanizing for free. Ring the surgery on 40541755 to fix a time to bring the birds in. Others also helping are Ulysses Vets at Stratford (40533977) and Orange Vets at Smithfield (40383283). Again, please phone and fix a time. Most other Vets will also euthanize for you. However, please contact to confirm.

For further advice email [cnsrim@gmail.com](mailto:cnsrim@gmail.com) or visit our website [www.cnsrim.org.au](http://www.cnsrim.org.au)